

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for    [Advanced Search](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display AbstractPlus Show 20 Sort By Send to

All: 1 Free Full Text: 0 Review: 0

1: [Circulation](#). 1989 Sep;80(3 Pt 1):I24-42.

[Links](#)

**Global and regional left ventricular systolic performance in the in situ ejecting canine heart. Importance of the mitral apparatus.**

[Sarris GE](#), [Fann JI](#), [Niczyporuk MA](#), [Derby GC](#), [Handen CE](#), [Miller DC](#).

Department of Cardiovascular Surgery, Stanford University School of Medicine, California 94305.

The importance of the intact mitral apparatus in left ventricular (LV) systolic performance has been indirectly suggested by clinical studies of chordal-preserving mitral valve replacement (MVR) or reconstruction. The importance of the intact mitral apparatus has been clearly demonstrated in isovolumic experimental preparations but has not been demonstrated unequivocally in ejecting hearts. Therefore, this question was assessed independently of load in an in situ, open-chest ejecting canine heart preparation (n = 9). Three orthogonal LV dimensions were measured by sonomicrometry. During MVR with complete chordal preservation, snares were placed around the anterior and posterior papillary muscles. After the hearts were weaned from cardiopulmonary bypass, LV function was assessed by caval occlusion to alter LV preload abruptly. The slopes of the end-systolic--pressure-volume (end-systolic elastance, Ees) and stroke-work--end-diastolic volume (preload-recrutable stroke work, PRSW) relations were used to measure global LV systolic function; similarly, the slopes of the end-systolic--pressure-dimension (regional end-systolic elastance, rEes) and stroke-work--end-diastolic dimension changes in regional LV systolic performance. All chordae were then divided by pulling the snares. Immediate reassessment revealed deterioration of global LV function: Ees declined by 72% (14.1 +/- 11.2 mm Hg/ml [mean +/- SD] vs. 3.9 +/- 3.5 mm Hg/ml, p less than 0.001), and PRSW declined by 39% (129 +/- 37 vs. 79 +/- 29 mm Hg, p = 0.0001). Regional LV function was also adversely affected but somewhat selectively: rEes decreased significantly in all three LV dimensions (p less than or equal to 0.03), but rPRSW decreased significantly (-21%) only in the anteroposterior minor LV axis (89 +/- 19 vs. 70 +/- 15 mm Hg, p = 0.005) and in the septal-lateral axis (-19%, p = NS). These data demonstrate the importance of the intact mitral apparatus on LV systolic performance in ejecting hearts, particularly in the LV regions subtended by the papillary muscles.

PMID: 2766532 [PubMed - indexed for MEDLINE]



**Related articles**

- Importance of mitral subvalvular apparatus in terms of cardiac energetics and systolic mech[J Clin Invest. 1991]
  - Mitral valve replacement in dilated canine hearts with chronic mitral regurgitation. Importanc [Circulation. 1991]
  - Importance of the mitral subvalvular apparatus for left ventricular segmental systolic mechar [Circulation. 1990]
  - [Review](#) Valvular-ventricular interaction: the importance of the mitral chordae tendineae in te[J Card Surg. 1988]
  - [Review](#) Mechanical loads and the isovolumic and filling indices of left ventricular rel; [Prog Cardiovasc Dis. 1990]
- » See reviews... | » See all...

**Cited by 1 PubMed Central article**

The effect of pure mitral regurgitation on mitral annular geometry and three-di [J Thorac Cardiovasc Surg. 2008]

**Recent Activity**

[Turn Off](#) [Clear](#)

- Global and regional left ventricular systolic performance in the in situ ejecting canine h...
- Surgical management of acute aortic dissection complicated by stroke.
- Role of the mitral subvalvular apparatus in left ventricular systolic mechanics.
- The reversibility of canine vein-graft arterialization.
- Treatment of patients with aortic dissection presenting with peripheral vascular complicat...

» See more...

Display AbstractPlus Show 20 Sort By Send to